

Chiral symmetry breaking beyond BCS and theorem on the width of scalars

P. Bicudo

Dep. Física and CFTP Instituto Superior Técnico

Abstract. We review chiral symmetry breaking at the BCS level, in the framework of chiral invariant quark models and in the Schwinger-Dyson formalism. We revisit the π mass problem beyond the BCS level. We show a theorem on the masses, on the widths and on the $q\bar{q}$ content of the scalar mesons σ and f_0 .

Keywords: Chiral Symmetry, BCS, Pion mass, Decay width

PACS: 11.30.Rd, 12.38.Lg, 13.25.Jx, 14.40.Aq

INTRODUCTION: CHIRAL SYMMETRY BREAKING AT BCS

With no quark loop, at the BCS level, equivalent to the rainbow/ladder truncation several hadronic properties can be derived in the quark model with many-body techniques pioneered by Scadron, and others [1, 2, 3, 4, 5] ,

- equivalent to the Schwinger-Dyson approach,
- and in the fermion perspective, equivalent to quenched Lattice QCD.

In this talk the equations are written in terms of Feynman diagrams, for simplicity. The propagators are quark propagators. The potential is not detailed since different quark-quark interactions are used in the literature. The interaction is only required

- to be chiral invariant, and this is sufficient to reproduce qualitatively the phenomena of low energy hadronic physics,
- to account quantitatively for the observables, say $M_\pi, f_\pi, M_\rho, \dots$.

At this BCS level we can start by computing the quark mass, solving the mass gap equation,

$$\text{---} \overleftarrow{\textcolor{red}{1}} \text{---}^{-1} = \text{---} \overleftarrow{\textcolor{red}{0}} \text{---}^{-1} - \text{---} \overleftarrow{\textcolor{red}{1}} \text{---}^{\text{wavy loop}}, \quad (1)$$

where 0 labels the free quark propagator and 1 labels the quark propagator at the BCS level, or ladder/rainbow truncation, in the series of Feynman diagrams.

Then, at the BCS level, the bound state equation is obtained summing the ladder,

$$\text{[Diagram: Square box with 1]} = \text{[Diagram: Two quarks connected]} + \text{[Diagram: Two quarks connected with one gluon exchange]} + \text{[Diagram: Two quarks connected with two gluon exchanges]} + \dots$$

$$\text{[Diagram: Square box with 1]} \approx \sum_M \text{[Diagram: Two quarks connected by } G_M \text{ with vertices } \Gamma_M]$$
(2)

For the vertex we get the Bethe-Salpeter equation,

$$\text{[Diagram: Vertex with two quarks and a dashed line]} = \text{[Diagram: Two quarks connected by a gluon exchange meeting at a vertex with a dashed line]}$$
(3)

which in the equal-time approximation is equivalent to the Salpeter or RPA equation. In any case one obtains the Gell-Mann-Oakes-Renner relations, implying that $M_\pi \rightarrow 0$ when $m_c \rightarrow 0$; and one also gets all the hadron spectrum, correct up to the quality of the potential,

$$\text{[Diagram: Quark-antiquark pair connected by a gluon line]}$$
(4)

The $\pi - \pi$ scattering can also be studied at the BCS level. The sum of all consistent series of ladder with one-loop diagrams sum was first performed by Hiller *et al.*, and by Boronowsky *et al.* [6, 7] in the NJL model. Then Bicudo, Llanes-Estrada, Cotanch *et al.* [8, 9, 10, 11] summed the same class of diagrams both in the quark model and in the Schwinger-Dyson formalism,

$$\text{[Diagram 1]} + \text{[Diagram 2]} - \text{[Diagram 3]}$$
(5)

and this indeed reproduces the Chiral theorems, say the Adler Zero and the Weinberg Theorem.

REVISITING THE π MASS PROBLEM BEYOND BCS

The BCS, or rainbow/ladder truncation, does not include coupled channels, nor decay widths of mesons. A correct description of hadronic physics requires the inclusion of coupled channel diagrams in the BS equation.

In effective field theories, without quarks or gluons, at least meson loops should be included,

$$\text{---}\pi\text{---} + \text{---}\pi\text{---} \text{---}\pi\text{---} + \text{---}\pi\text{---} \text{---}\pi\text{---} \text{---}\pi\text{---} + \text{---}\pi\text{---} \text{---}\pi\text{---} \text{---}\pi\text{---} + \text{---}\pi\text{---} \text{---}\pi\text{---} \text{---}\pi\text{---} \text{---}\pi\text{---} \quad (6)$$

Applying this to the pion, it would produce a negative shift of M_π^2 , or imaginary shift of M_π . Thus this problem must be solved self-consistently.

At the quark level we must supplement the Bethe Salpeter kernel for the pion (Isospin $I = 1$) with the diagrams,

Interestingly, this may account for the η' , η , π_0 mass differences, supplementing the isospin $I = 1$ diagrams with the diagrams that only contribute to the $I = 0$ mesons, as in

Lakhina and Bicudo [12], with the diagrams,

(8)

A THEOREM ON SCALAR σ , $f_0 \cdots$ WIDTHS AND $\langle q\bar{q} \rangle$ CONTENT


To again solve the M_π imaginary mass problem, now beyond the BCS level, we have to return to the mass gap equation, going beyond BCS, now with at least one extra quark loop.

(9)

where 2 labels the quark propagator beyond BCS, at 1-loop level.

This contribution to the mass gap equation was computed by Bicudo [13], see also Wojciech Broniowski *et al* [7].

Importantly, the new beyond BCS term suppresses chiral symmetry breaking, like in the σ model the ϕ^4 term counteracts the $-\phi^2$ term,



(10)

and this term must necessarily be smaller than the BCS level terms, to preserve the spontaneous breaking of chiral symmetry.

Notice that this term increases with the decay width of the isospin $I = 0$ scalars with a large $\langle q\bar{q} \rangle$ component, and with the mass of these same $I = 0$ scalars. Thus we arrive at a theorem stating that,

- either the isospin $I = 0$ scalars have a small decay
- or they have a large mass
- or they have a small $\langle q\bar{q} \rangle$ component.

SUMMARY

To conclude we find that,

- either the $q\bar{q}$ coupling to the scalar meson is small,
- or the scalar meson mass M is large,
- or the $\pi\pi$ coupling to the scalar meson is small.

Moreover the chiral invariant quark model or equivalently the Schwinger-Dyson formalism reproduce several chiral theorems, and also reproduces the $U_A(1)$ splittings of the $M_{\eta'}$, M_{η} , M_{π_0} masses.

The results in a particular confining and chiral invariant quark model [13] are,

- this one loop beyond BCS rainbow/ladder truncation decreases the quark condensate $\langle q\bar{q} \rangle$ by 5% to 50%.
- the lowest $q\bar{q}$ scalar has a mass $M \simeq 1$ GeV, noticing however that the spin-orbit interaction is too strong and thus a better model might produce a larger mass,
- the lowest $q\bar{q}$ scalar has a width Γ between 50 MeV and 100 MeV.

REFERENCES

1. Y. Nambu and G. Jona-Lasinio, "Dynamical Model Of Elementary Particles Based On An Analogy With Superconductivity. I," Phys. Rev. **122**, 345 (1961); Y. Nambu and G. Jona-Lasinio, "Dynamical Model Of Elementary Particles Based On An Analogy With Superconductivity. II," Phys. Rev. **124**, 246 (1961).
2. R. Delbourgo and M. D. Scadron, "Proof Of The Nambu-Goldstone Realization For Vector Gluon Quark Theories," J. Phys. G **5**, 1621 (1979).
3. A. Le Yaouanc, L. Oliver, S. Ono, O. Pene and J. C. Raynal, "A Quark Model Of Light Mesons With Dynamically Broken Chiral Symmetry," Phys. Rev. D **31**, 137 (1985).
4. P. Bicudo and J. E. Ribeiro, "Current Quark Model In A P Wave Triplet Condensed Vacuum. 1. The Dynamical Breaking Of Chiral Symmetry," Phys. Rev. D **42**, 1611 (1990); Phys. Rev. D **42**, 1625; Phys. Rev. D **42**, 1635.
5. F. J. Llanes-Estrada and S. R. Cotanch, "Meson structure in a relativistic many-body approach," Phys. Rev. Lett. **84**, 1102 (2000) [arXiv:hep-ph/9906359].
6. V. Bernard, U. G. Meissner, A. Blin and B. Hiller, "Four point functions in quark flavor dynamics: Meson meson scattering," Phys. Lett. B **253**, 443 (1991).
7. E. N. Nikolov, W. Broniowski, C. V. Christov, G. Ripka and K. Goeke, Nucl. Phys. A **608**, 411 (1996) [arXiv:hep-ph/9602274].
8. P. Bicudo, "Analytic proof that the quark model complies with partially conserved axial current theorems," Phys. Rev. C **67**, 035201 (2003).
9. P. Bicudo, S. Cotanch, F. Llanes-Estrada, P. Maris, J. E. Ribeiro and A. Szczepaniak, "Chirally symmetric quark description of low energy pi pi scattering," Phys. Rev. D **65**, 076008 (2002) [arXiv:hep-ph/0112015].
10. P. Bicudo, M. Faria, G. M. Marques and J. E. Ribeiro, "Microscopic derivation of N (Delta) pi scattering lengths," Nucl. Phys. A **735**, 138 (2004) [arXiv:nucl-th/0106071].
11. F. J. Llanes-Estrada and P. Bicudo, "Quark Schwinger-Dyson evaluation of the l_1 , l_2 coefficients in the chiral Lagrangian," Phys. Rev. D **68**, 094014 (2003) [arXiv:hep-ph/0306146].
12. O. Lakhina and P. Bicudo, "Microscopic quark study of the eta and eta' masses," arXiv:0708.0984 [hep-ph].
13. P. Bicudo, "New perspective on the scalar meson puzzle, from spontaneous chiral symmetry breaking beyond BCS," Phys. Rev. C **60**, 035209 (1999).